

①⑫

**EUROPEAN PATENT APPLICATION**

①⑮ Application number: **88310000.0**

①⑮ Int. Cl.4: **A61K 37/02 , C12N 15/00 ,**  
**C12N 1/18 , C12P 21/02 ,**  
**//C12N5/00**

①⑯ Date of filing: **25.10.88**

The microorganism(s) has (have) been deposited  
with The National Collection of Industrial and  
Marine Bacteria under number NCIB 12242

The title of the invention has been amended  
(Guidelines for Examination in the EPO, A-III,  
7.3).

①⑳ Priority: **30.10.87 GB 8725529**

④③ Date of publication of application:  
**28.06.89 Bulletin 89/26**

⑧④ Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI LU NL SE**

⑦① Applicant: **Delta Biotechnology Limited**  
**Castle Court, Castle Boulevard**  
**Nottingham NG7 1FD(GB)**

⑦② Inventor: **Ballance, David James**  
**11 South Road**  
**West Bridgford Nottingham NG2 7AG(GB)**  
Inventor: **Hinchliffe, Edward**  
**16 Lambley Lane**  
**Burton Joyce Nottingham NG14 5BG(GB)**  
Inventor: **Geisow, Michael John**  
**115 Main Street**  
**East Bridgford Nottingham NG13 8NH(GB)**  
Inventor: **Senior, Peter James**  
**Wilson House Wilson**  
**Near Melbourne Derbyshire(GB)**

⑦④ Representative: **Bassett, Richard Simon et al**  
**ERIC POTTER & CLARKSON 14 Oxford Street**  
**Nottingham NG1 5BP(GB)**

⑤④ **N-terminal fragments of human serum albumin.**

⑤⑦ Polypeptides corresponding to mature human serum albumin residues 1 to n, where n is between 369 and 419 inclusive, are useful as substitutes for albumin in the treatment of burns and shock in humans, the clearances of undesirable compounds, (such as bilirubin) from human blood, in laboratory growth media and in HSA assays.

**EP 0 322 094 A1**

HSA (1-389) is particularly preferred, although not novel per se.  
The polypeptides may be produced by recombinant DNA techniques, especially in yeast.

## POLYPEPTIDES

This invention relates to a novel polypeptide molecule which can be produced by recombinant DNA technology and can be used for many of the existing applications of human serum albumin.

Human serum albumin (HSA) is the most abundant plasma protein, contributing 60% w/w of the total protein content of the plasma. A molecule of HSA consists of a single non-glycosylated polypeptide chain of  
 5 585 amino acids of formula molecular weight 66,500. The amino acid sequence of HSA has been established by protein sequence analysis (Meloun et al, 1975, "Complete amino acid sequence of human serum albumin" FEBS. Letters: 58:1, 136-317; Behrens et al, 1975, "Structure of human serum albumin" Fed. Proc. 34, 591) and more recently by genetic analysis (Lawn et al, 1981, Nucleic Acids Research 9, 6102-6114). Although there have been discrepancies between the amino acid sequences as published  
 10 (some being attributable to polymorphisms), Figure 1 represents the amino acid sequence currently believed to be most representative of the HSA present within the human population.

Because of its relatively small molecular weight and net negative charge at physiological pH (Peters, 1970, "Serum albumin", Adv. Clin. Chem. 13, 37-111), HSA contributes 85% of the osmotic effect of normal plasma. Thus HSA is the principal regulator of plasma volume. A secondary role of HSA is to bind small  
 15 molecules produced by catabolic processes (for example fatty acids and bilirubin). Albumin represents the principal means for the transport of these key metabolites, which are poorly soluble at physiological pH. Physical, chemical, immunological and limited proteolytic studies of HSA have shown that the molecule is composed of regions of polypeptide chains which retain their conformation after separation from the parent molecule by enzymatic means. These polypeptide chains retain their binding capabilities thereby facilitating  
 20 the mapping of binding sites for bilirubin, fatty acids and other small molecules to particular regions of the polypeptide chain (Kragh-Hansen, 1981, "Molecular aspects of ligand binding to serum albumin". A. Soc. Pharm. Expt. Ther. 33, 1, 17-53). Much of the information in this area has been reviewed (Brown and Shockey, 1982, "Serum albumin: structure and characterisation of its ligand binding sites").

The indications for the clinical use of therapeutic concentrates of HSA are related principally to its  
 25 oncotic action as a plasma volume expander. Concentrates of HSA have been used therapeutically since the 1940's, in particular in cases of shock, burns, adult respiratory distress syndrome, and cardiopulmonary bypass. Albumin has also been used in cases of acute liver failure, following removal of ascitic fluid from patients with cirrhosis, after surgery, in acute nephrosis, in renal dialysis, and as a transport protein for removing toxic substances, such as in severe jaundice in haemolytic disease of the new born.

30 In addition to its use as a therapeutic agent, HSA is a major component of serum added to media used to support the growth of mammalian cells in tissue culture. The consumption of serum and hence of albumin has been greatly increased over recent years as biotechnology and pharmaceutical companies have expanded their tissue culture for research and for production. There is a universal need for lower cost and better regulation of sera for these purposes.

35 It is known to manipulate the HSA-encoding DNA sequence express a recombinant polypeptide in microorganisms. Indeed such a recombinant HSA polypeptide has been produced in bacterial species such as *Escherichia coli* (G.B. Patent No. 2 147 903B) and *Bacillus subtilis* (European Patent Application No. 86304656.1) and the yeast *Saccharomyces cerevisiae* (European Patent Publication No. 201 239, Delta Biotechnology Ltd.); thus it is generally accepted that a recombinant polypeptide essentially identical to  
 40 natural HSA can be produced in a variety of microbial hosts by employing known methods. However, in all cases where recombinant HSA has been produced, the objective has been to produce a molecule which is "nature-identical" to HSA in structure and biological function.

It has now been found that it is advantageous to produce shorter forms of HSA.

One aspect of the present invention provides a polypeptide comprising the N-terminal portion of human  
 45 serum albumin up to amino acid residue n, where n is 369 to 419, and variants thereof.

The novel polypeptides of the invention are hereinafter referred to as "HSA(1-n)".

The terms "human serum albumin" is intended to include (but not necessarily to be restricted to) known or yet-to-be discovered polymorphic forms of HSA. For example, albumin Naskapi has Lys-372 in place of Glu-372 and pro-albumin Christchurch has an altered pro-sequence. The term "variants" is  
 50 intended to include (but not necessarily to be restricted to) minor artificial variations in residues 1 to n (such as molecules lacking one or a few residues, having conservative substitutions or minor insertions of residues, or having minor variations of amino acid structure). Thus polypeptides which have 80%, preferably 85%, 90%, 95% or 99%, homology with any HSA (1-n) compound are deemed to be "variants". Such variants are preferably 360 to 430 amino acids long, more preferably 369 to 419 amino acids long and most preferably 386 to 388 amino acids long. It is also preferred for such variants to be physiologically equivalent

to HSA (1-n) compounds; that is to say, variants preferably share at least one pharmacological utility with HSA (1-n) compounds. Furthermore, any putative variant which is to be used pharmacologically should be non-immunogenic in the animal (especially human) being treated.

Conservative substitutions are those where one or more amino acids are substituted others having similar properties such that one skilled in the art of polypeptide chemistry would expect at least the secondary structure, and preferably the tertiary structure, of the polypeptide to be substantially unchanged. For example, typical such substitutions include alanine or valine for glycine, arginine or asparagine for glutamine, serine for asparagine and histidine for lysine. Variants may alternatively, or as well, lack up to ten (preferably only one or two) amino acid residues in comparison with any given HSA (1-n); preferably any such omissions occur in the 100 to 369 portion of the molecule (relative to mature HSA itself). Similarly, up to ten, but preferably only one or two, amino acids may be added, again in the 100 to 369 portion for preference. The term "physiologically functional equivalents" also encompasses larger molecules comprising the said 1 to n sequence plus a further sequence at the N-terminal (for example, pro-HSA(1-n), pre-pro-HSA(1-n), met-HSA(1-n), and HSA(1-n) having a suitable leader sequence which is not necessarily native to HSA).

If the HSA (1-n) is to be prepared by culturing a transformed yeast (*S. cerevisiae*) as is described in more detail below, the leader sequence may, for example, be that found naturally with the yeast alpha-factor protein. C-terminal fusion products with other polypeptides of interest may be produced. Known forms and fragments of HSA are clearly to be regarded as excluded from the above definition, for example HSA(1-387), which was a peptic fragment produced in low yield (Geisow and Beaven, *Biochem. J.* **161**, 619-624, 1977 and *ibid.* **163**, 477-484, 1977. These prior articles identify the fragment as 1-386, but it has since become apparent (see, for example, Lawn *et al.*, *op.cit.*) that this is due to the authors' use of incorrect published sequence information and that the fragment was in fact 1-387). Similarly, a C-terminal fusion protein comprising HSA (1-n) and the remaining HSA residues (numbers n + 1 to 585) is not claimed as part of the invention.

Particularly preferred novel HSA(1-n) compounds include HSA(1-373) (i.e. C-terminal Val), HSA(1-388) (i.e. C-terminal Ile), HSA(1-389) (i.e. C-terminal Lys), HSA(1-390) (i.e. C-terminal Gln) and HSA(1-407) (i.e. C-terminal Leu).

The HSA(1-n) molecules are preferably produced by means for recombinant DNA technology (optionally followed by proteolytic digestion), rather than by chemical or enzymatic degradation of natural HSA, or by peptide synthesis. In the case of enzymatic degradation, for example, a trypsin-like enzyme will cleave HSA between Lys(389) and Gln(390) but also concomitantly at other cleavage sites. In the future, peptide synthesis may become more feasible for molecules as long as 419 amino acids, but at present is not a practical proposition. Expression in yeast is particularly preferred.

It has been found that, at least in some situations where the HSA(1-n) compound is produced by culturing a transformed host, some HSA(1-n) compounds which are longer than HSA(1-387) are proteolytically digested back to HSA (1-387) by the enzymes which are naturally present in the system. Thus, one can, if desired, use a nucleotide sequence corresponding to a given HSA(1-n) compound in order to prepare another HSA(1-n) compound.

The new molecules herein described can be used as an effective substitute for either natural HSA or nature-identical recombinant HSA as a plasma volume expander. An advantage of HSA(1-n) over natural HSA and recombinant nature-identical HSA relates to the efficacy of raising the colloid osmotic pressure of blood. The smaller molecular weight (approximately 44 kilo-daltons) of the protein of the present invention means that an individual protein dose of only one-half to two-thirds that of natural-HSA or nature-identical recombinant HSA will be required for the equivalent colloid osmotic effect. Consequently, any process for the production of this novel polypeptide by means of recombinant DNA technology may afford significant economic advantages over known processes for the production of nature-identical recombinant HSA, since substantially less proteinaceous material is required to be produced for an effective dose.

Thus, a second aspect of the invention provides a pharmaceutical composition comprising HSA(1-n)-plus, where HSA(1-n)plus is HSA(1-n) as defined above or any HSA(1-n) molecules which are known per se but have not been proposed for pharmaceutical use.

HSA (1-387) which, as discussed above, was a fragment produced by chance in a prior art peptic digest of HSA, is particularly preferred as the HSA(1-n) plus in such a pharmaceutical composition. The composition may comprise "variants" of HSA (1-387) as defined above.

A third aspect provides a method of treating a human for shock, burns or other conditions in which albumin is indicated, comprising administering intravenously a blood-bulking or blood-clearing effective non-toxic amount of a sterile non-pyrogenic solution of a polypeptide comprising HSA(1-n) plus.

Further aspects of the invention include (a) vectors, plasmids and transformed microorganisms,

including cell lines, encoding HSA(1-n)plus expression; (b) processes for the production of HSA(1-n)plus comprising the fermentation under suitable conditions of a microorganism (including a cell line) so transformed as to express HSA(1-n)plus; and (c) laboratory media comprising HSA(1-n)plus.

A further advantage of at least some HSA(1-n) plus molecules over nature-identical recombinant HSA is that their smaller size and thus reduced amino acid content has been found to lead to an increase in the yield obtained (molecules per cell dry weight) in microbial hosts relative to that obtained currently for nature-identical recombinant HSA. Thus, not only has it been found that the scale of the process can be reduced, but also productivity in the recombinant host organism can be enhanced.

The compounds of the invention may be used as blood-bulking (plasma-expanding) agents in analogous ways and in analogous formulations as HSA itself except that the dose of the HSA(1-n)plus compound (in terms of weight) will generally be less than that of HSA as the oncotic effect of the former is greater. The pharmacist or clinician skilled in the art will readily be able to determine by routine and non-inventive experimentation the optimum dose of the HSA(1-n)plus compound. Generally, the amount of HSA(1-n)plus which is administered will be about two-thirds of the amount of HSA which would be administered.

HSA (1-n) plus compounds may also be used as:  
(1) substitutes for HSA or, more commonly, bovine serum albumin (BSA) in tissue culture media, thereby reducing the risk of contamination of the medium with, for example, viruses and mycoplasmas; (2) substitutes for BSA in the stationary phase in liquid chromatography for resolution of enantiomers and so on.

### EXAMPLES

The invention will now be illustrated by way of example and with reference to the drawings, in which:

Figure 1 depicts the amino acid sequence currently thought to be the most representative of natural HSA, with (boxed) the alternative C-terminal of HSA(1-n);

Figure 2 depicts the DNA sequence coding for mature HSA;

Figure 3 illustrates, diagrammatically, the construction of mHOB16;

Figure 4 illustrates, diagrammatically, the construction of pHOB31; and

Figure 5 is a copy of a rocket electrophoretogram showing the increased yield of HSA(1-389) over complete HSA.

Standard recombinant DNA procedures are as described by Maniatis *et al* (1982) unless otherwise stated. Construction and analysis of M13 recombinant clones was as described by Messing (1983) and Sanger *et al.* (1977).

The human serum albumin coding sequence used in the construction of the following molecules is derived from the plasmid M13mp19.7 (European Patent Application No. 201 239, Delta Biotechnology Ltd.) or by synthesis of oligonucleotides equivalent to parts of this sequence. Oligonucleotides were synthesised using phosphoramidite chemistry on an Applied Biosystems 380B oligonucleotide synthesizer according to the manufacturer's recommendations (AB Inc., Warrington, Cheshire, England).

#### Example 1: HSA (1-389)

An expression vector was constructed in which DNA encoding the HSA secretion signal and mature HSA up to and including the 389th amino acid, lysine, was placed downstream of the *S.cerevisiae* phosphoglycerate kinase gene (PGK) promoter and followed by a stop codon and the PGK terminator of transcription. This vector was then introduced into *S.cerevisiae* by transformation and directed the expression and secretion from the cells of a molecule representing the N-terminal 389 amino acids of HSA.

An oligonucleotide was synthesised (Linker 1) which represented a part of the known HSA coding sequence (Figure 2) from the PstI site (1092, Figure 2) to the codon for valine 381 wherein that codon was changed from GTG to GTC:

Linker 1

D P H E C Y A K V F D E  
 5' GAT CCT CAT GAA TGC TAT GCC AAA GTG TTC GAT GAA  
 3' ACGT CTA GGA GTA CTT ACG ATA CGG TTT CAC AAG CTA CTT  
 5 1100 1120  
 F K P L V  
 10 TTT AAA CCT CTT GTC 3'  
 AAA TTT GGA GAA CAG 5'

Linker 1 was ligated into the vector M13mp19 (Norrander et al, 1983) which had been digested with PstI and HincII and the ligation mixture was used to transfect E.coli strain XL1-Blue (Stratagene Cloning Systems, San Diego, CA). Recombinant clones were identified by their failure to evolve a blue colour on medium containing the chromogenic indicator X-gal (5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactoside) in the presence of IPTG (isopropylthio- $\beta$ -galactoside). DNA sequence analysis of template DNA prepared from bacteriophage particles of recombinant clones identified a molecule with the required DNA sequence, designated mHOB12 (Figure 3).

M13mp19.7 consists of the coding region of mature HSA in M13mp19 (Norrander et al, 1983) such that the codon for the first amino acid of HSA, GAT, overlaps a unique XhoI site thus:

25 Asp Ala  
 5' C T C G A G A T G C A 3'  
 3' G A G C T C T A C G T 5'  
 XhoI

(EPA No. 210239 A1). M13mp19.7 was digested with XhoI, made flush-ended by S1-nuclease treatment and was then ligated with the following oligonucleotide (Linker 2):

35 Linker 2

40 5' T C T T T T A T C C A A G C T T G G A T A A A G A 3'  
 3' A G A A A A T A G G T T C G A A C C T A T T T T C T 5'  
 HindIII

45 The ligation mix was then used to transfect E.coli XL1-Blue and template DNA was prepared from several plaques and then analysed by DNA sequencing to identify a clone, pDBD1 (Figure 4), with the correct sequence.

A 1.1 kb HindIII to PstI fragment representing the 5' end of the HSA coding region and one half of the inserted oligonucleotide linker was isolated from pDBD1 by agarose gel electrophoresis. This fragment was then ligated with double stranded mHOB12 previously digested with HindIII and PstI and the ligation mix was then used to transfect E.coli XL1-Blue. Single stranded template DNA was prepared from mature bacteriophage particles of several plaques. The DNA was made double stranded *in vitro* by extension from annealed sequencing primer with the Klenow fragment of DNA polymerase I in the presence of deoxynucleoside triphosphates. Restriction enzyme analysis of this DNA permitted the identification of a clone with the correct configuration, mHOB15 (Figure 4).

The following oligonucleotide (Linker 3) represents from the codon for the 382nd amino acids of mature HSA (glutamate, GAA) to the codon for lysine 389 which is followed by a stop codon (TAA) and a HindIII site and then a BamHI cohesive end:

Linker 3

5                    E   E   P   Q   N   L   I   K   J  
 5'       GAA GAG CCT CAG AAT TTA ATC AAA TAA GCTTG       3'  
 3'       CTT CTC GGA GTC TTA AAT TAG TTT ATT CGAACCTAG       5'

10       This was ligated into double stranded mHOB15, previously digested with HincII and BamHI. After ligation, the DNA was digested with HincII to destroy all non-recombinant molecules and then used to transfect E.coli XL1-Blue. Single stranded DNA was prepared from bacteriophage particles of a number of clones and subjected to DNA sequence analysis. One clone having the correct DNA sequence was designated mHOB16 (Figure 4).

15       A molecule in which the mature HSA coding region was fused to the HSA secretion signal was created by insertion of Linker 4:

Linker 4

20                    M   K   W   V   S   F   I   S   L   L   F   L  
 5'   GATCC ATG AAG TGG GTA AGC TTT ATT TCC CTT CTT TTT CTC  
 25       G TAC TCC ACC CAT TCG AAA TAA AGG GAA GAA AAA GAG  
  
 F   S   S   A   Y   S   R   G   V   F   R   R  
 30       TTT AGC TCG GCT TAT TCC AGG GGT GTG TTT CG       3'  
 AAA ACG AGC CGA ATA AGG TCC CCA CAC AAA GCAGCT       5'

35       into BAMHI and XhoI digested M13mp19.7 to form pDBD2 (Figure 5). In this linker the codon for the fourth amino acid after the initial methionine, ACC for threonine in the HSA pre-pro leader sequence (Lawn et al, 1981), has been changed to AGC for serine to create a HindIII site.

40       The 5' end of this construction was removed as a BamHI to PvuII fragment and ligated with the PvuII to BamHI fragment of double stranded mHOB16 (representing the 3' end of the truncated HSA gene) into pMA91 (Mellor et al, 1983) at the BglII site to form pHOB31 (Figure 4). This molecule contains the truncated HSA coding region with the HSA secretion signal between the S.cerevisiae PGK gene promoter and terminator such that the 5' end of the gene abuts the promoter. The molecule also contains a selectable marker for yeast transformation, LEU2, and part of the yeast 2um plasmid to permit autonomous replication in yeast.

45       The plasmid pHOB31 was introduced into S.cerevisiae AH22 (Hinnen et al, 1978) by transformation using standard procedures (Beggs, 1978). Purified transformants were grown in YEPD broth (1% yeast extract, 2% peptone, 2% glucose) for 3 days at 30°C and the culture supernatant was then analysed, successfully, for the presence of HSA-related material by rocket gel electrophoresis. Figure 5 shows the electrophoretogram; the yield of HSA-related material from transformants harbouring a plasmid encoding HSA(1-389) is demonstrably higher than the yield from a transformant secreting mature, natural, HSA.

50       However, production of HSA (1-389) gave a product indistinguishable from HSA (1-387) (see Example 2) by both amino-terminal and carboxy-terminal sequence analysis. This is probably explained by the efficient removal of the COOH-terminal sequence Ile-Lys.

EXAMPLE 2: HSA (1-387)

55       The construction of a plasmid encoding HSA (1-387) was identical to the procedure for construction of the HSA (1-389) plasmid, pHOB31, except that the linker 3 was substituted by linker 5 (shown below) which represents the region from the codon for the 382nd amino acid of mature HSA (glutamate, GAA) to the

codon for leucine 387 which is followed by a stop codon and a HindIII site and then a BamHI cohesive end:

Linker 5

5

```

          E   E   P   Q   N   L   Stop
5'  GAA GAG CCT CAG AAT TTA TAA GCTTG      3'
10 3'  CTT CTC GGA GTC TTA AAT ATT CGAACCTAG 5'

```

The remainder of the construction was as detailed above for pHOB31 and resulted in the plasmid pDBD5.

15 EXAMPLE 3: (1-369)

In order to construct a plasmid encoding HSA (1-369), a linker was synthesised representing the region from the PstI site of mature HSA (position 1092, Figure 3) to the codon for cystine 369 which was followed by a stop codon (TAA), a HindIII site and then a BamHI cohesive end:

20

Linker 6

```

          D   P   H   E   C   Stop
25 5'          GAT  CCT  CAT  GAA  TGC  TAA GCTTG
    3'  A  CGT  CTA  GGA  GTA  CTT  ACG  ATT CGAACCTAG

```

30 This linker was ligated with the BamHI PstI fragment of pDBD2, representing the 5' part of preproHSA, into pMA91 at the BglII site. A plasmid with the correct configuration was termed pDBD3 (Figure 6).

Production of HSA (1-369) by culturing S.cerevisiae transformed with pDBD3 gave low yields, indicating that the product may have been unstable in the yeast expression system used.

35

EXAMPLE 4: HSA (1-419)

40 For the construction of a plasmid encoding HSA (1-419) the BamHI - HincII fragment of pDBD2 was ligated with an annealed self-complementary oligonucleotide (linker 7):

Linker 7

45 5' ATAAGCTTGGATCCAAGCTTAT 3'

and then the ligation mix was digested with BamHI and the fragment was ligated into pMA91 to give pDBD4 (Figure 7). In this construct the HincII site (1256, Figure 3) of pDBD2 creates a blunt end after the second base of the codon for serine 419 and this codon is reformed by the linker 6 such that this codon is followed by a stop codon, a HindIII site and a BamHI site.

50 Expression of HSA (1-419) via plasmid pDBD5 in S.cerevisiae produced a molecule with the correct amino terminal sequence (Asp-Ala-His.....) but leucine and not serine was the COOH-terminal residue. Attempts to isolate the COOH-terminal peptide using a covalent label which should attach to cysteine 392 also were unsuccessful. It was concluded that proteolysis of part of the COOH-terminus of HSA (1-419) occurred. This is consistent with the observation of a small percentage of proteolysis in the same position of  
55 full-length HSA produced in an analogous manner in yeast (Sleep et al. 1988).

EXAMPLE 5: Fermentation of HSA(1-n)plus-producing yeast

- A laboratory fermenter is filled to half its nominal working volume with an initial "batch" medium containing 50ml/l of a salts mixture (containing 114g/l  $\text{KH}_2\text{PO}_4$ , 12g/l  $\text{MgSO}_4$ , 3.0g/l  $\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$ , 2.0g/l  $\text{Na}_2\text{EDTA}$ , 10ml/l of a trace elements solution containing 3g/l  $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ , 10g/l  $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ , 3.2g/l  $\text{MnSO}_4 \cdot 4\text{H}_2\text{O}$ , 79mg/l  $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ , 1.5g/l  $\text{H}_3\text{BO}_3$ , 0.2g/l  $\text{KI}$ , 0.5g/l  $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ , 0.56g/l  $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ , 75ml/l  $\text{H}_3\text{PO}_4$ : 20g/l sucrose: 50ml/l of a vitamins mixture containing 1.6g/l Ca pantothenate, 1.2g/l nicotinic acid, 12.8g/l m inositol, 0.32g/l thiamine HCl and 8mg/l pyridoxine HCl and 8mg/l biotin. An equal volume of "feed" medium containing 100ml/l of the salts mixture, 20ml/l of trace elements solution 500g/l sucrose and 100ml/l vitamin solution is held in a separate reservoir connected to the fermenter by a metering pump.
- 10 The fermenter is inoculated with *Saccharomyces cerevisiae* which has been transformed as above with plasmid pDBD3 from Example 2. The pH is maintained at  $5.7 \pm 0.2$  by automatic addition of ammonia or sulphuric acid, the temperature is kept at  $30^\circ\text{C}$  and the stirred speed is adjusted to give a dissolved oxygen tension (DOT) of  $> 20\%$  air saturation at 1 v/v/min air flow rate. When the initial substrate has been consumed, the metering pump is turned on, maintaining a growth rate of approximately  $0.15\text{h}^{-1}$ . The pump
- 15 rate is increased to maintain this growth rate until the stirrer speed reached its maximum value at which point it is not possible to increase the pump rate any further without causing the DOT to fall below 15% air saturation which is the minimum value permitted to occur. PPG 2000 is added in response to a foam sensor. None is added until over 50% of the feed solution had been added. The final level of addition is 0.2g/l.
- 20 HSA(1-387) is secreted into the medium

#### EXAMPLE 6: Binding of bilirubin to HSA(1-387)

- 25 Binding of the haem metabolite, bilirubin, to HSA (1-387) was carried out by a fluorescence enhancement method (Beaven and Gratzen (1973) Eur. J. Biochem. 33, 500-510). Figure 8 shows that the enhancement of bilirubin fluorescence as a function of protein/bilirubin ratio is indistinguishable for HSA(1-387) and clinical grade HSA.
- The interaction of HSA and bilirubin is very sensitive to the conformation of the protein (Beaven and Gratzen, loc. cit.) and these results indicate that no gross alteration in conformation of the regions of HSA
- 30 represented by HSA(1-387) has occurred through the expression of a shorter molecule.

#### EXAMPLE 7: Oncotic behaviour of HSA(1-387)

- 35 HSA(1-387) was concentrated in 0.9% w/v saline to a final protein concentration of 54 mg/ml. Dilutions of this concentrate, together with dilutions of a clinical grade HSA (100 mg/ml), were compared for osmotic effect in a colloid osmometer. Figure 9 indicates that HSA(1-387) gives a colloid osmotic pressure approximately one-third higher than that of full-length HSA at a given protein concentration. Importantly, the
- 40 increase in colloid osmotic pressure with protein concentration is approximately linear over a range up to 5% w/v, which represents the concentration in plasma.
- This indicates that HSA(1-387) does not self-associate appreciably within a useful working clinical concentration range.

#### EXAMPLE 8: Formulations for Injection

- 45 The HSA(1-n)plus of the invention may be presented in container sizes ranging from 20ml to 500ml, with the concentration thereof varying (typically) from 2% to 17%, for example 3%, 13% or 17%.
- 50 The solution for administration is sterile and pyrogen free. A 3% solution is osmotically similar to human plasma. At least 96% of the total protein is preferably albumin. The sodium ion content is generally between 130-160mmol/litre and the potassium ion content is generally not more than 2mmol/litre. The pH is adjusted to  $6.9 \pm 0.5$ . The concentration of citrate is generally no more than 20mmol/litre and may be absent altogether.
- 55 Stabilizers may be used, for example either 0.16 millimole sodium acetyl tryptophanate, or 0.08 millimole sodium acetyl tryptophanate and 0.08 millimole sodium caprylate per gram of HSA(1-n)plus.



## References

- Beggs, J.D. (1978). *Nature*. 275, 104-109.
- Brown, J.R. and Shockley, P., (1982) in "Lipid-Protein Interactions" 1, 25-68. Eds. Hayes, O. and Jost, P.C.
- 5 Hinnen, A. et al (1978). *Proc. Natl. Acad. Sci. USA*, 75, 1929-1933.
- Lawn, R.M. et al (1981). *Nucl. Acid. Res.* 9, 6103-6114.
- Maniatis, T. et al (1982). *Molecular cloning: A laboratory manual*. Cold Spring Harbour Laboratory, Cold Spring Harbor, New York.
- Mellor, J. et al (1983). *Gene*, 24, 1-14.
- 10 Messing, J. (1983). *Methods Enzymol.* 101, 20-78.
- Norlander, J. et al (1983). *Gene*, 26, 101-106.
- Sanger, F. et al (1977). *Proc. Natl. Acad. Sci. USA*, 74, 5463-5467.
- Sleep, D. Belfield, G.P. and Goodey, A.R. (1988) *Yeast* 4, S168.

15

## Claims

1. A polypeptide comprising the N-terminal portion of mature human serum albumin up to amino acid residue n, where n is 369 to 419, but not 387, and variants thereof.
- 20 2. A polypeptide according to Claim 1 wherein the polypeptide is selected from the group consisting of HSA (1-373), HSA (1-388), HSA (1-389), HSA (1-390) and HSA (1-407) and variants thereof.
3. A pharmaceutical composition comprising a polypeptide according to Claim 1 or 2 except that n may be 387.
4. A composition according to Claim 3 wherein the polypeptide is HSA (1-387) or a variant thereof.
- 25 5. A nucleotide sequence encoding a polypeptide comprising the N-terminal portion of mature human serum albumin up to amino acid residue n, where n is 309 to 419, and polypeptide variants thereof, the nucleotide sequence not being linked at its 3' end to a further sequence encoding the C-terminal portion of mature human serum albumin from amino acid residue n + 1 to 585.
6. A nucleotide sequence according to Claim 5 wherein n is 387.
- 30 7. A nucleotide sequence according to Claim 5 or 6 linked at its 5' end to a further nucleotide sequence encoding a peptide corresponding to the pro-, pre-, or pre-pro- position of HSA, a methionine residue, or another leader sequence.
8. An expression vector suitable for transformation of and expression in a selected host, the vector comprising a nucleotide sequence according to any one of Claims 6 and 8 and the said nucleotide
- 35 sequence being a DNA sequence.
9. A host organism transformed with a vector according to Claim 8.
10. A host organism according to Claim 9 which is Saccharomyces cerevisiae.
11. A process for the production of a polypeptide comprising the culture under suitable conditions of a host microorganism according to Claim 9 or 10, the said polypeptide being encoded by the said nucleotide
- 40 sequence.
12. A laboratory medium for the growth of microorganisms comprising a polypeptide according to Claim 1, except that n may be 387.
13. A medium according to Claim 12 wherein n is 387.

45

50

55

FIGURE 1

	10		20
Asp	Ala	His	Lys
Ser	Glu	Val	Ala
His	Arg	Phe	Lys
Asp	Leu	Gly	Glu
Glu	Asn	Phe	Lys
	30		40
Ala	Leu	Val	Leu
Ile	Ala	Phe	Ala
Gln	Tyr	Leu	Gln
Gln	Cys	Pro	Phe
Glu	Asp	His	Val
	50		60
Lys	Leu	Val	Asn
Glu	Val	Thr	Glu
Phe	Ala	Lys	Thr
Cys	Val	Ala	Asp
Glu	Ser	Ala	Glu
	70		80
Asn	Cys	Asp	Lys
Ser	Leu	His	Thr
Leu	Phe	Gly	Asp
Lys	Leu	Cys	Thr
Val	Ala	Thr	Leu
	90		100
Arg	Glu	Thr	Tyr
Gly	Glu	Met	Ala
Asp	Cys	Cys	Ala
Lys	Gln	Glu	Pro
Glu	Pro	Glu	Arg
Asn	Glu		
	110		120
Cys	Phe	Leu	Gln
His	Lys	Asp	Asp
Asn	Pro	Asn	Leu
Pro	Arg	Leu	Val
Arg	Pro	Glu	Val
	130		140
Asp	Val	Met	Cys
Thr	Ala	Phe	His
Asp	Asn	Glu	Glu
Thr	Phe	Leu	Lys
Lys	Lys	Tyr	Leu
Tyr			
	150		160
Glu	Ile	Ala	Arg
Arg	His	Pro	Tyr
Phe	Tyr	Ala	Pro
Glu	Leu	Leu	Phe
Phe	Ala	Lys	Arg
	170		180
Tyr	Lys	Ala	Ala
Phe	Thr	Glu	Cys
Cys	Gln	Ala	Ala
Asp	Lys	Ala	Ala
Cys	Leu	Leu	Pro
	190		200
Lys	Leu	Asp	Glu
Leu	Arg	Asp	Glu
Gly	Lys	Ala	Ser
Ser	Ala	Lys	Gln
Arg	Leu	Lys	Cys
	210		220
Ala	Ser	Leu	Gln
Lys	Phe	Gly	Glu
Arg	Ala	Phe	Lys
Ala	Trp	Ala	Val
Ala	Arg	Leu	Ser
	230		240
Gln	Arg	Phe	Pro
Lys	Ala	Glu	Phe
Ala	Glu	Val	Ser
Lys	Leu	Val	Thr
Asp	Leu	Thr	Lys
	250		260
Val	His	Thr	Glu
Cys	Cys	His	Gly
Asp	Leu	Leu	Glu
Cys	Ala	Asp	Asp
Arg	Ala	Asp	Leu
	270		280
Ala	Lys	Tyr	Ile
Cys	Glu	Asn	Gln
Asp	Ser	Ile	Ser
Ser	Lys	Leu	Lys
Glu	Cys	Cys	Glu
	290		300
Lys	Pro	Leu	Leu
Glu	Lys	Ser	His
Cys	Ile	Ala	Glu
Val	Glu	Asn	Asp
Glu	Met	Pro	Ala
	310		320
Asp	Leu	Pro	Ser
Leu	Ala	Ala	Asp
Phe	Val	Glu	Ser
Lys	Asp	Val	Cys
Lys	Asn	Tyr	Ala
	330		340
Glu	Ala	Lys	Asp
Val	Phe	Leu	Gly
Met	Phe	Leu	Tyr
Glu	Tyr	Ala	Arg
Arg	His	Pro	Asp
	350		360
Tyr	Ser	Val	Val
Leu	Leu	Leu	Arg
Leu	Ala	Lys	Thr
Tyr	Glu	Thr	Thr
Leu	Glu	Lys	Cys
	370		380
Cys	Ala	Ala	Ala
Asp	Pro	His	Glu
Cys	Tyr	Ala	Lys
Val	Phe	Asp	Glu
Phe	Lys	Pro	Leu

FIGURE 1 Cont.

Val Glu Glu Pro Gln Asn Leu Ile Lys	390	Gln Asn Cys Glu Leu Phe Glu Gln Leu Gly Glu	400
Tyr Lys Phe Gln Asn Ala Leu Leu Val	410	Arg Tyr Thr Lys Lys Val Pro Gln Val Ser Thr	420
Pro Thr Leu Val Glu Val Ser Arg Asn	430	Leu Gly Lys Val Gly Ser Lys Cys Cys Lys His	440
Pro Glu Ala Lys Arg Met Pro Cys Ala	450	Glu Asp Tyr Leu Ser Val Val Leu Asn Gln Leu	460
Cys Val Leu His Glu Lys Thr Pro Val	470	Ser Asp Arg Val Thr Lys Cys Cys Thr Glu Ser	480
Leu Val Asn Arg Arg Pro Cys Phe Ser	490	Ala Leu Glu Val Asp Glu Thr Tyr Val Pro Lys	500
Glu Phe Asn Ala Glu Thr Phe Thr Phe	510	His Ala Asp Ile Cys Thr Leu Ser Glu Lys Glu	520
Arg Gln Ile Lys Lys Gln Thr Ala Leu	530	Val Glu Leu Val Lys His Lys Pro Lys Ala Thr	540
Lys Glu Gln Leu Lys Ala Val Met Asp	550	Asp Phe Ala Ala Phe Val Glu Lys Cys Cys Lys	560
Ala Asp Asp Lys Glu Thr Cys Phe Ala	570	Glu Glu Gly Lys Lys Leu Val Ala Ala Ser Gln	580
Ala Ala Leu Gly Leu			

FIGURE 2 DNA sequence coding for mature HSA

10 20 30 40 50 60 70 80  
GATGCACACAAGAGTGAGGTTGCTCATCGGTTTAAAGATTTGGGAGAAGAAAATTTCAAAGCCTTGGTGTGATTGCCTT  
D A H K S E V A H R F K D L G E E N F K A L V L I A F

90 100 110 120 130 140 150 160  
TGCTCAGTATCTTCAGCAGTGTCCATTTGAAGATCATGTAAAATTAGTGAATGAAGTAACTGAATTTGCAAAAACATGTG  
A Q Y L Q Q C P F E D H V K L V N E V T E F A K T C

170 180 190 200 210 220 230 240  
TTGCTGATGAGTCAGCTGAAAATTGTGACAAATCACTTCATACCCTTTTTGGAGACAAATTATGCACAGTTGCAACTCTT  
V A D E S A E N C D K S L H T L F G D K L C T V A T L

250 260 270 280 290 300 310 320  
CGTGAAACCTATGGTGAATGGCTGACTGCTGTGCAAAACAAGAACCTGAGAGAAATGAATGCTTCTTGCAACACAAAGA  
R E T Y G E M A D C C A K Q E P E R N E C F L Q H K D

330 340 350 360 370 380 390 400  
TGACAACCCAAACCTCCCCGATTGGTGAGACCAGAGGTTGATGTGATGTGCACTGCTTTTCATGACAATGAAGAGACAT  
D N P N L P R L V R P E V D V M C T A F H D N E E T

410 420 430 440 450 460 470 480  
TTTTGAAAAATACTTATATGAAATTGCCAGAAGACATCCTTACTTTTATGCCCGGAACTCCTTTTCTTGCTAAAAGG  
F L K K Y L Y E I A R R H P Y F Y A P E L L F F A K R

490 500 510 520 530 540 550 560  
TATAAAGCTGCTTTTACAGAATGTGCCAAGCTGCTGATAAAGCTGCCTGCCTGTTGCCAAAGCTCGATGAACCTTCGGGA  
Y K A A F T E C C Q A A D K A A C L L P K L D E L R D

570 580 590 600 610 620 630 640  
TGAAGGGAAGGCTTCGTCTGCCAAACAGAGACTCAAATGTGCCAGTCTCCAAAAATTTGGAGAAAGAGCTTTCAAAGCAT  
E G K A S S A K Q R L K C A S L Q K F G E R A F K A

650 660 670 680 690 700 710 720  
GGGAGTGGCTCGCCTGAGCCAGAGATTTCCCAAAGCTGAGTTTGCAGAAGTTTCCAAGTTAGTGACAGATCTTACAAA  
W A V A R L S Q R F P K A E F A E V S K L V T D L T K

730 740 750 760 770 780 790 800  
GTCCACACGGAATGCTGCCATGGAGATCTGCTTGAATGTGCTGATGACAGGGCGGACCTTGCCAAGTATATCTGTGAAAA  
V T E C C H G D L L E C A D D R A D L A K Y I C E N

810 820 830 840 850 860 870 880  
TCAGGATTCGATCTCCAGTAAACTGAAGGAATGCTGTGAAAAACCTCTGTTGGAAAAATCCCACTGCATTGCCGAAGTGG  
Q D S I S S K L K E C C E K P L L E K S H C I A E V

890 900 910 920 930 940 950 960  
AAAATGATGAGATGCCTGCTGACTTGCCCTTCATTAGCTGCTGATTTTGTGAAAGTAAGGATGTTTGCAAAAACATATGCT  
E N D E M P A D L P S L A A D F V E S K D V C K N Y A

970 980 990 1000 1010 1020 1030 1040  
GAGGCAAAGGATGTCTTCTGGGCATGTTTTGTATGAATATGCAAGAAGGCATCCTGATTACTCTGTCGTGCTGCTGCT  
E A K D V F L G M F L Y E Y A R R H P D Y S V V L L L

FIGURE 2 Cont.

1050 1060 1070 1080 1090 1100 1110 1120  
GAGACTTGCCAAGACATATGAAACCACTCTAGAGAAGTGCTGTGCCGCTGCAGATCCTCATGAATGCTATGCCAAAGTGT  
R L A K T Y E T T L E K C C A A A D P H E C Y A K V

1130 1140 1150 1160 1170 1180 1190 1200  
TCGATGAATTTAAACCTCTTGTGGAAGAGCCTCAGAATTTAATCAAACAAAACCTGTGAGCTTTTTGAGCAGCTTGGAGAG  
F D E F K P L V E E P Q N L I K Q N C E L F E Q L G E

1210 1220 1230 1240 1250 1260 1270 1280  
TACAAATTCAGAATGCGCTATTAGTTCGTTACACCAAGAAAGTACCCCAAGTGTCAACTCCAACCTCTGTAGAGGTCTC  
Y K F Q N A L L V R Y T K K V P Q V S T P T L V E V S

1290 1300 1310 1320 1330 1340 1350 1360  
AAGAAACCTAGGAAAAGTGGGCAGCAAATGTTGTAAACATCCTGAAGCAAAAAGAATGCCCTGTGCAGAAGACTATCTAT  
R N L G K V G S K C C K H P E A K R M P C A E D Y L

1370 1380 1390 1400 1410 1420 1430 1440  
CCGTGGTCTGAACCAAGTTATGTGTGTTGCATGAGAAAACGCCAGTAAGTGACAGAGTCACAAAATGCTGCACAGAGTCC  
S V V L N Q L C V L H E K T P V S D R V T K C C T E S

1450 1460 1470 1480 1490 1500 1510 1520  
TTGGTGAACAGGCGACCATGCTTTTCAGCTCTGGAAGTCGATGAAACATACGTTCCCAAAGAGTTTAATGCTGAAACATT  
L V N R R P C F S A L E V D E T Y V P K E F N A E T F

1530 1540 1550 1560 1570 1580 1590 1600  
CACCTTCCATGCAGATATATGCACACTTTCTGAGAAGGAGAGACAAATCAAGAAACAACTGCACTTGTTGAGCTTGTGA  
T F H A D I C T L S E K E R Q I K K Q T A L V E L V

1610 1620 1630 1640 1650 1660 1670 1680  
AACACAAGCCCAAGGCAACAAAAGAGCAACTGAAAGCTGTTATGGATGATTTTCGCAGCTTTTGTAGAGAAGTGCTGCAAG  
K H K P K A T K E Q L K A V M D D F A A F V E K C C K

1690 1700 1710 1720 1730 1740 1750 1760  
GCTGACGATAAGGAGACCTGCTTTGCCGAGGAGGGTAAAAAACTTGTTGCTGCAAGTCAAGCTGCCTTAGGCTTATAACA  
A D D K E T C F A E E G K K L V A A S Q A A L G L

1770 1780  
TCTACATTTAAAAGCATCTCAG

FIGURE 3 Construction of mHOB16




FIGURE 4 Construction of pHOB31






FIGURE 5

Rocket immunoelectrophoretic analysis of culture supernatant from S.cerevisiae AH22 transformants obtained with a plasmid containing the complete HSA coding region (samples 1-4) and from transformants harbouring an equivalent plasmid encoding truncated HSA (1-389) (samples 5-8).




FIGURE 6 Construction of pDBD3



Le meilleur est déposé


**FIGURE 7**



Neu cingericht / Newy filed  
Nouvellement / Newy filed


FIGURE 8

Titration of bilirubin by BPL HSA and  $\Delta 21$   
(measurement of fluorescence enhancement)



Neu eingereicht / Newly filed  
Nouvellement déposé

FIGURE 9





DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X,D	THE BIOCHEMICAL JOURNAL, vol. 163, no. 3, 1977, pages 477-484, GB; M.J. GEISOW et al.: "Physical and binding properties of large fragments of human serum albumin" * Abstract * ---	4	A 61 K 37/02 C 12 N 15/00 C 12 N 1/18 C 12 P 21/02 // C 12 N 5/00
A	THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 257, no. 6, 25th March 1982, pages 2770-2774, US; N. DOYEN et al.: "Immunochemical cross-reactivity between cyanogen bromide fragments of human serum albumin" * Page 2770, column 2, paragraph 1 * ---	1-9	
A	EP-A-0 073 646 (GENENTECH INC.) * Figure 2; page 12, lines 16-18 * ---	1-9,11	
A	EP-A-0 201 239 (DELTA BIOTECHNOLOGY LTD) * Figure 3; example II * -----	1-11	
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
			C 12 N C 12 P
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 09-02-1989	Examiner CUPIDO M.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons ..... & : member of the same patent family, corresponding document	